PHASE ENHANCED ABSORPTION

A Novel Method To Enhance Absorption Rate

DOE Awarding Number: DE-FG26-04NT42128

(2004.7 - 2007.6)

Principal Author: Dr. Liang Hu

Students: Victor Roberts, Jr.

Monica J. Wood

School of Engineering and Technology Hampton University Hampton, VA 23668

DOE Project Manager: Tim Four

INTRODUCTION

Physical absorption

the gas component being absorbed is simply dissolved in the liquid absorbent.

CO₂ dissolves into Water, Propylene Carbonate, Methanol, etc.

Chemical Absorption
 there is a chemical reaction between the gas component being absorbed and a component in the liquid to form a compound.

Hot Potassium Carbonate Solution, Aqueous Alkanolamines, etc.

Chemical Absorption

- Advantage
 - 1) enhancing the absorption rate
 - 2) increasing carrying capacities for gas components
- Disadvantage
 chemical reaction
 causes difficulty for
 the release of the gas
 components from
 liquid

What is Phase Enhanced Absorption?

In phase enhanced absorption, more than one liquid phases were involved in the absorption process. One of the liquid phase was called as carrying phase, in which gas solute will be finally accumulated. Another liquid phase is known as transportation phase. The transportation phase only plays the rule for the transportation of the gas solute from gas phase to carrying phase and for the increase of absorption rate. In phase enhanced absorption, the absorption rate is enhanced by the transportation phase.

What is Phase Enhanced Absorption?

Gas


Aqueous solution

Gas

Organic layer

Aqueous solution

MASS TRANSFER MODEL



MASS TRANSFER EQUATION

Physical Absorption

$$N_A = k_1 (C_A^* - C_A)$$

Chemical Absorption

$$N_A = E_R k_1 (C_A^* - C_A)$$

Phase Enhanced Absorption

$$N_A = E_P k_1 (C_A^* - C_A)$$

STIRRING CELL




FIGURE 1 STIRRING CELL

EXPERIMENTAL FLOW CHART





FIGURE 2 EXPERIMENTAL APPARATUS


- $1 \texttt{CO2} \ \texttt{CYLINDER}; \quad 2 \texttt{REGULATOR}; \quad 3 \texttt{BUFFEL} \ \texttt{BOTTLE}; \quad 4 \texttt{PRESSURE} \ \texttt{STABLE} \ \texttt{TUBE};$
- 5,8 ROTATING FLOW METER; 6 SILICON GEL U TUBE; 7 ACTIVE CARBON U TUBE;
- 9-SATURATOR; 10,12-FOAM FILM FLOW METER; 11-MOTER; 13-LIQUID FEED FUNNEL;
- 14 STIRRING CELL: 15 THERMOSTAT: T1. T2. T3 THERMOMETER

RESULTS AND DISCUSSION


 Following experimental results prove that CO₂ absorption rate is enhanced by PHASE


CO₂-water system and CO₂-water-n-haptane layer system






CO₂-water system and CO₂-water-isooctanol system






CO_2 -potassium carbonate aqueous solution CO_2 -potassium carbonate aqueous solution-organic layer (membrane 3 mm)



${\rm CO_2}$ -potassium carbonate aqueous solution ${\rm CO_2}$ -potassium carbonate aqueous solution-organic layer (membrane 9 mm)




MECHANISM

- Mass transfer resistant between the interface of gas and liquid is much higher than that between the interface of liquid and liquid
- Mass transfer resistant between the interface of gas and organic layer is lower than that between the interface of gas and aqueous solution
- The transportation layer (organic layer) delivers CO₂ from gas phase into water phase and enhances the absorption rate as long as the sufficient mass transfer (agitation) between two liquid phases is provided.

CONCLUSION

- Organic layer can increase the absorption rate as long as sufficient mass transfer in liquid (agitation) is provided
- Organic layer plays the role of transportation of CO₂ from gas to aqueous solution
- However, the enhanced factor is highly depended on the liquid mass transfer coefficient (agitation)

APPLICATION



WHAT IS NEXT?

• Seeking the application of phase enhanced absorption in CO₂ separation from flue gas